



Clarus™

PRO

PROFESSIONAL LANDSCAPE FERTILIZER

Nada Lotta Yada

16-0-3

WITH 1% IRON

MEDIUM SIZE

GUARANTEED ANALYSIS

Total Nitrogen (N).....	16.0%
3.15% Ammoniacal Nitrogen	
12.32% Urea Nitrogen*	
0.53% Water Insoluble Nitrogen**	
Soluble Potash (K ₂ O).....	3.0%
Calcium (Ca)	1.0%
Sulfur (S)	4.0%
4.0% Combined Sulfur (S)	
Iron (Fe).....	1.0%

PLANT NUTRIENTS DERIVED FROM

Poultry Manure, Biosolids, Ammonium Sulfate, Sulfur Coated Urea, Urea, Sulfate of Potash

*4.74 slowly available urea nitrogen from sulfur coated urea.

**0.53% slowly available water insoluble nitrogen from poultry manure and biosolids.

F1519

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

NET WEIGHT
50 LB (22.7 kg)

DISTRIBUTED & GUARANTEED BY

Nutrients PLUS, LLC
2133 Upton Dr, Ste 126-509 • VA Beach, VA 23454
(888) 675-7515 • www.nutrientsplus.com

Nutrients PLUS, LLC warrants that this product conforms to the analysis on its label. When used in accordance with label directions, under normal conditions, this product is reasonably fit for its intended purposes. Since time, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

FOR MORE INFORMATION SEE REVERSE SIDE

Nada Lotta Yada

16-0-3

WITH 1% IRON

MEDIUM SIZE

STORAGE: Store in original container in a cool, dry place. Do not contaminate food or feedstuffs by storage and disposal.

TURF 16-0-3 can be applied to any type of grass at any time of year with no risk of burn when used as directed. It is always good practice to water in fertilizer materials. Cool Season Grasses: Apply in spring and twice during the fall. Warm Season Grasses: Apply every two months during the growing season. Prior to Seeding or Sodding: Apply to area to be seeded or sodded.

COVERAGE

	<u>LOW RATE</u>	<u>RECOMMENDED RATE</u>	<u>HIGH RATE</u>
Coverage in ft ² of turf	16,000	11,400	8,000
	For .5 lbs. of <u>N/1000 ft² of turf</u>	For .7 lbs. of <u>N/1000 ft² of turf</u>	For 1.0 lbs. of <u>N/1000 ft² of turf</u>
Lbs per 1000 ft ² of turf	3.125	4.375	6.25
Lbs per acre	136	190	272
Lbs K per 1000 ft ² of turf	0.094	0.13	0.188
	Delivers .25 lbs. of Organic Matter per 1000 ft ²	Delivers .35 lbs. of Organic Matter per 1000 ft ²	Delivers .5 lbs. of Organic Matter per 1000 ft ²

RECOMMENDED RATE SPREADER SETTINGS

MEDIUM SIZE

EARTHWAY ROTARY	13
LESCO	F/G
PRIZELAWN BF-1	I
SCOTTS R8	K
SPYKER/CYCLONE	5
LELY	7
VICON	28

Spreader Settings Are Approximate, You May Need To Adjust For Proper Coverage.

FLORIDA APPLICATIONS: For use on golf courses or other athletic turf in FL, we recommend you follow SL191, "Recommendations for N, P, K & Mg for Golf Course and Athletic Field Fertilization Base on Mehlich I Extractant", available on the internet at <http://edis.ifas.ufl.edu/SS404>. For commercial application to urban turf or lawns in FL, we recommend you follow the Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries, available on the internet at <http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP Book final.pdf>.